

## New England Biolabs Certificate of Analysis

*Product Name:* Q5<sup>®</sup> High-Fidelity DNA Polymerase  
*Catalog #:* M0491S/L  
*Concentration:* 2,000 units/ml  
*Unit Definition:* One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 74°C  
*Lot #:* 0051612  
*Assay Date:* 12/2016  
*Expiration Date:* 12/2018  
*Storage Temp:* -20°C  
*Storage Conditions:* Proprietary  
*Specification Version:* PS-M0491S/L v2.0  
*Effective Date:* 31 Mar 2016


Assay Name/Specification (minimum release criteria)	Lot #0051612
<b>Endonuclease Activity (Nicking, Polymerase)</b> - A 50 µl reaction in NEBuffer 2 in the presence of 400 µM dNTPs containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 units of Q5 <sup>®</sup> High-Fidelity DNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>PCR Amplification (20 kb Lambda DNA)</b> - A 50 µl reaction in Q5 <sup>®</sup> Reaction Buffer in the presence of 200 µM dNTPs and 1.0 µM primers containing 10 ng Lambda DNA with 1 unit of Q5 <sup>®</sup> High-Fidelity DNA Polymerase for 22 cycles of PCR amplification results in the expected 20 kb product.	<b>Pass</b>
<b>PCR Amplification (7 kb Human Genomic DNA)</b> - A 50 µl reaction in Q5 <sup>®</sup> Reaction Buffer in the presence of 200 µM dNTPs and 0.5 µM primers containing 20 ng Human Genomic DNA with 1 unit of Q5 <sup>®</sup> High-Fidelity DNA Polymerase for 30 cycles of PCR amplification results in the expected 7 kb product.	<b>Pass</b>
<b>PCR Amplification (Enhancer Dependent, &gt;65% GC-rich)</b> - A 50 µl reaction in Q5 <sup>®</sup> Reaction Buffer and Q5 <sup>®</sup> High GC Enhancer in the presence of 200 µM dNTPs and 0.5 µM primers containing 20 ng Human Genomic DNA with 1 unit of Q5 <sup>®</sup> High-Fidelity DNA Polymerase for 30 cycles of PCR amplification results in the enhancer-dependent production of the expected 452 bp product.	<b>Pass</b>
<b>Phosphatase Activity (pNPP)</b> - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl <sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Q5 <sup>®</sup> High-Fidelity DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	<b>Pass</b>

## New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0051612
<p><b>Protein Purity Assay (SDS-PAGE)</b> - Q5<sup>®</sup> High-Fidelity DNA Polymerase is <math>\geq 95\%</math> pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>qPCR DNA Contamination (<i>E. coli</i> Genomic)</b> - A minimum of 2 units of Q5<sup>®</sup> High-Fidelity DNA Polymerase is screened for the presence of <i>E. coli</i> genomic DNA using SYBR<sup>®</sup> Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is <math>\leq 1</math> <i>E. coli</i> genome.</p>	<b>Pass</b>
<p><b>RNase Activity (Extended Digestion)</b> - A 10 <math>\mu</math>l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 <math>\mu</math>l of Q5<sup>®</sup> High-Fidelity DNA Polymerase is incubated at 37°C. After incubation for 16 hours, <math>&gt;90\%</math> of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<b>Pass</b>



Authorized by  
Melanie Fortier  
31 Mar 2016



Inspected by  
Tony Spear-Alfonso  
16 Dec 2016

