

New England Biolabs Certificate of Analysis

Product Name: PNGase F
Catalog Number: P0704S
Concentration: 500,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10 µg of denatured RNase B in 1 hour at 37°C in a total reaction volume of 10 µl (65 NEB units = 1 IUB milliunit).
Packaging Lot Number: 10231463
Expiration Date: 01/2026
Storage Temperature: -20°C
Storage Conditions: 50 mM NaCl , 20 mM Tris-HCl , 5 mM EDTA , 50 % Glycerol, (pH 7.5 @ 25°C)
Specification Version: PS-P0704S/L v1.0

PNGase F Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0704SVIAL	PNGase F	10226044	Pass
B3704SVIAL	10X GlycoBuffer 2	10227056	Pass
B2704SVIAL	NP-40	10203694	Pass
B1704SVIAL	Glycoprotein Denaturing Buffer	10227027	Pass

Assay Name/Specification	Lot # 10231463
Endoglycosidase F1 Activity (LC/MS) A 20 µl reaction in Glyco Buffer 2 containing 20 pmoles of 2-AA Man-5 fluorescent standard and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no endoglycosidase F1 activity as determined by LC/MS analysis with fluorescent detection.	Pass
Glycosidase Activity (Endo F1, F2, H) A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (Endo F2, F3) A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 5,000 units of PNGase F	Pass

Assay Name/Specification	Lot # 10231463
<p>incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (α-Glucosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fucα1-2)Galβ1-4Glc-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Neuraminidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-2 Fucosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucα1-2Galβ1-4Glc-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Fucosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucα1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Galactosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Mannosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-3Manβ1-4GlcNAc-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by</p>	Pass

Assay Name/Specification	Lot # 10231463
thin layer chromatography.	
<p>Glycosidase Activity (α1-6 Galactosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Mannosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Mannosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNACβ1-4Galβ1-4Glc-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNACβ1-4GlcNACβ1-4GlcNAC-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Xylosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNACβ1-4Galβ1-4Glc-AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as</p>	Pass

Assay Name/Specification	Lot # 10231463
determined by thin layer chromatography.	
<p>Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC) and 5,000 units of PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Protease Activity (SDS-PAGE) A 20 μl reaction in 1X Glyco Buffer 2 containing 24 μg of a standard mixture of proteins and a minimum of 10,000 units of PNGase F incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) PNGase F is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Maxwell Elkus
Production Scientist
07 Feb 2024



Michael Tonello
Packaging Quality Control Inspector
21 Mar 2024