

## New England Biolabs Certificate of Analysis

**Product Name:** Endo F3  
**Catalog Number:** P0771S  
**Concentration:** 8,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to cleave > 95% of the carbohydrate from 10 µg Porcine Fibrinogen in 1 hour at 37°C in a total reaction volume of 10 µl.  
**Packaging Lot Number:** 10152120  
**Expiration Date:** 06/2024  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)  
**Specification Version:** PS-P0771S/L v1.0

Endo F3 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0771SVIAL	Endo F3	10152119	Pass
B1703SVIAL	10X Glycobuffer 4	10134776	Pass

Assay Name/Specification	Lot # 10152120
<p><b>Glycosidase Activity (Endo F1, F2, H)</b>            A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p><b>Protein Purity Assay (SDS-PAGE)</b>            Endo F3 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p><b>Protease Activity (SDS-PAGE)</b>            A 20 µl reaction in 1X Glyco Buffer 4 containing 24 µg of a standard mixture of proteins and a minimum of 80 units of Endo F3 incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	Pass
<p><b>Glycosidase Activity (α1-6 Mannosidase)</b>            A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 40 units of Endo F3</p>	Pass

Assay Name/Specification	Lot # 10152120
<p>incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p><b>Glycosidase Activity (α1-6 Galactosidase)</b> A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (α1-3 Mannosidase)</b> A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-3Manβ1-4GlcNAc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (α1-3 Galactosidase)</b> A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (α1-3 Fucosidase)</b> A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (α1-2 Fucosidase)</b> A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-2Galβ1-4Glc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β-Xylosidase)</b> A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β-N-Acetylglucosaminidase)</b> A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as</p>	<b>Pass</b>

Assay Name/Specification	Lot # 10152120
determined by thin layer chromatography.	
<p><b>Glycosidase Activity (<math>\beta</math>-N-Acetylgalactosaminidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled <math>\beta</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\beta</math>-Mannosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled <math>\beta</math>-Mannosidase substrate (Man<math>\beta</math>1-4Man<math>\beta</math>1-4Man-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\beta</math>1-4 Galactosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled <math>\beta</math>-Galactosidase substrate (Gal<math>\beta</math>1-4GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\beta</math>1-3 Galactosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled <math>\beta</math>-Galactosidase substrate (Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-Neuraminidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Neuraminidase substrate (Neu5Ac<math>\alpha</math>2-3Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-N-Acetylgalactosaminidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 4 containing 1 mM of fluorescently-labeled <math>\alpha</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\alpha</math>1-3(Fuc<math>\alpha</math>1-2)Gal<math>\beta</math>1-4Glc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-Glucosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Glucosidase substrate (Glc<math>\alpha</math>1-6Glc<math>\alpha</math>1-4Glc-AMC) and 40 units of Endo F3 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



---

Alicia Bielik  
Production Scientist  
15 Jun 2022



---

Erin Varney  
Packaging Quality Control Inspector  
15 Jun 2022