

New England Biolabs Certificate of Analysis

Product Name: Endo F2
Catalog #: P0772S
Concentration: 8,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to cleave >95% of the carbohydrate from 10 µg Porcine Fibrinogen in 1 hour at 37°C in a total reaction volume of 10 µl.
Lot #: 0011710
Assay Date: 10/2017
Expiration Date: 10/2019
Storage Temp: -20°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0772S v1.0
Effective Date: 11 Nov 2016

Assay Name/Specification (minimum release criteria)	Lot #0011710
Glycosidase Activity (β-Mannosidase) - A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-Xylosidase) - A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-3 Galactosidase) - A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-4 Galactosidase) - A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylgalactosaminidase) - A 10 µl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0011710
Glycosidase Activity (β-N-Acetylglucosaminidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β -N-Acetylglucosaminidase substrate (GlcNAc β 1-4GlcNAc β 1-4GlcNAc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-Glucosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Glucosidase substrate (Glc α 1-6Glc α 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-Neuraminidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Neuraminidase substrate (Neu5Ac α 2-3Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-2 Fucosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuca1-2Gal β 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Fucosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuca1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Galactosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-3Gal β 1-4GlcNAc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Mannosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-3Man β 1-4GlcNAc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-6 Galactosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-6Gal α 1-6Glc α 1-2Fru-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-6 Mannosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-6Man α 1-6(Man α 1-3)Man-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-N-Acetylgalactosaminidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -N-Acetylgalactosaminidase substrate (GalNAc α 1-3(Fuca1-2)Gal β 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass



New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0011710
Protease Activity (SDS-PAGE) - A 20 µl reaction in 1X Glyco Buffer 4 containing 24 µg of a standard mixture of proteins and a minimum of 40 units of Endo F2 incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue.	Pass
Protein Purity Assay (SDS-PAGE) - Endo F2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass



Authorized by
Derek Robinson
11 Nov 2016



Inspected by
Brad Landgraf
03 Oct 2017

