

New England Biolabs Certificate of Analysis

Product Name: *HpaI*
Catalog Number: *R0105L*
Concentration: *5,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10173219*
Expiration Date: *12/2024*
Storage Temperature: *-20°C*
Storage Conditions: *10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @25°C)*
Specification Version: *PS-R0105S/L/V v2.0*


HpaI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0105LVIAL	HpaI	10173217	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10167588	Pass
B6004SVIAL	rCutSmart™ Buffer	10168650	Pass

Assay Name/Specification	Lot # 10173219
Protein Purity Assay (SDS-PAGE) HpaI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 15 units of HpaI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with HpaI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with HpaI.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of HpaI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results	Pass

Assay Name/Specification	Lot # 10173219
<p>are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p> <p>Endonuclease Activity (Nicking) A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of supercoiled pUC19 DNA and a minimum of 15 units of HpaI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p> <p>Exonuclease Activity (Radioactivity Release) A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of HpaI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	<p style="text-align: center;">Pass</p> <p style="text-align: center;">Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



YunJie Sun
Production Scientist
06 Dec 2022



Michael Tonello
Packaging Quality Control Inspector
07 Dec 2022