

New England Biolabs Certificate of Analysis

Product Name: *SacII*
Catalog Number: *R0157S*
Concentration: *20,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10056468*
Expiration Date: *06/2021*
Storage Temperature: *-20°C*
Storage Conditions: *50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA*
Specification Version: *PS-R0157S/L v1.0*

| SacII Component List | | | |
|----------------------|------------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R0157SVIAL | SacII | 10046227 | Pass |
| B7204SVIAL | CutSmart® Buffer | 10046090 | Pass |
| B7024SVIAL | Gel Loading Dye, Purple (6X) | 10046084 | Pass |

| Assay Name/Specification | Lot # 10056468 |
|---|----------------|
| <p>Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 60 Units of SacII incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.</p> | Pass |
| <p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of SacII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p> | Pass |
| <p>Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pXba DNA with SacII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SacII.</p> | Pass |
| <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 60 Units of SacII incubated for 16 hours at 37°C results in a DNA pattern free of</p> | Pass |

| Assay Name/Specification | Lot # 10056468 |
|---|----------------|
| detectable nuclease degradation as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.



Anthony Francis
Production Scientist
07 Jun 2019



Jay Minichiello
Packaging Quality Control Inspector
04 Nov 2019