

## New England Biolabs Certificate of Analysis


**Product Name:** *Bsrl*  
**Catalog Number:** *R0527S*  
**Concentration:** *10,000 U/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme required to digest 1 µg of PhiX174 DNA in 1 hour at 65°C in a total reaction volume of 50 µl.*  
**Packaging Lot Number:** *10233290*  
**Expiration Date:** *02/2026*  
**Storage Temperature:** *-20°C*  
**Storage Conditions:** *300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA*  
**Specification Version:** *PS-R0527S/L v2.0*

Bsrl Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0527SVIAL	Bsrl	10228348	Pass
B6003SVIAL	NEBuffer™ r3.1	10227733	Pass

Assay Name/Specification	Lot # 10233290
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 50 units of Bsrl incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of PhiX174 DNA with Bsrl, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with Bsrl.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of PhiX174 DNA and a minimum of 50 units of Bsrl incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



---

Ana Egana  
Production Scientist  
01 Mar 2024



---

Michael Tonello  
Packaging Quality Control Inspector  
01 Mar 2024