


## New England Biolabs Certificate of Analysis

**Product Name:** AclI  
**Catalog Number:** R0598S  
**Concentration:** 5,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot Number:** 10026011  
**Expiration Date:** 11/2020  
**Storage Temperature:** -20°C  
**Storage Conditions:** 100 mM KCl , 10 mM Tris-HCl (7.4), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 200 µg/ml BSA  
**Specification Version:** PS-R0598S/L v1.0


AclI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0598SVIAL	AclI	10026012	Pass
B7204SVIAL	CutSmart® Buffer	10021118	Pass

Assay Name/Specification	Lot # 10026011
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 15 units of AclI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with AclI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AclI.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> AclI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H ] E. coli DNA and a minimum of 50 units of AclI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

This product has been tested and shown to be in compliance with all specifications.



Tony Spear-Alfonso  
Production Scientist  
01 Oct 2018



Michael Tonello  
Packaging Quality Control Inspector  
31 Oct 2018