

## New England Biolabs Certificate of Analysis

**Product Name:** BbvCI  
**Catalog Number:** R0601S  
**Concentration:** 2,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart™ Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10182472  
**Expiration Date:** 03/2024  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml rAlbumin (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0601S/L v3.0


BbvCI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0601SVIAL	BbvCI	10182471	Pass
B6004SVIAL	rCutSmart™ Buffer	10182167	Pass

Assay Name/Specification	Lot # 10182472
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 10 units of BbvCI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 2-fold over-digestion of Lambda DNA with BbvCI, 95% can be recut with BbvCI.	Pass
<b>Non-Specific DNase Activity (16 hour)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 2 units of BbvCI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> BbvCI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue	Pass

Assay Name/Specification	Lot # 10182472
<p>detection.</p> <p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 2 units of BbvCI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is <math>\leq 1</math> E. coli genome.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



YunJie Sun  
Production Scientist  
20 Mar 2023



Michael Tonello  
Packaging Quality Control Inspector  
26 Apr 2023