

New England Biolabs Certificate of Analysis

Product Name: *SrfI*
Catalog Number: *R0629L*
Concentration: *20,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of pNEB193-SrfI DNA in CutSmart incubated for 1 hour at 37°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10182402*
Expiration Date: *09/2024*
Storage Temperature: *-20°C*
Storage Conditions: *300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)*
Specification Version: *PS-R0629S/L v1.0*

SrfI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0629LVIAL	SrfI	10182401	Pass
B6004SVIAL	rCutSmart™ Buffer	10176423	Pass

Assay Name/Specification	Lot # 10182402
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 100 units of SrfI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 200 units of SrfI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in CutSmart® Buffer containing 1 µg of pNEB193-SrfI DNA and 1 µl of SrfI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pNEB193-SrfI DNA with SrfI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	Pass

