

## New England Biolabs Certificate of Analysis

**Product Name:** SrfI  
**Catalog Number:** R0629S  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of pNEB193-SrfI DNA in CutSmart incubated for 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10122891  
**Expiration Date:** 03/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0629S/L v1.0

SrfI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0629SVIAL	SrfI	10122889	Pass
B6004SVIAL	rCutSmart™ Buffer	10124441	Pass

Assay Name/Specification	Lot # 10122891
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of pNEB193-SrfI DNA and a minimum of 20 units of SrfI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of pNEB193-SrfI DNA with SrfI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SrfI.	Pass
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of pNEB193-SrfI DNA and 1 µl of SrfI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and	Pass


Assay Name/Specification	Lot # 10122891
double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of SrfI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 100 units of SrfI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Protein Purity Assay (SDS-PAGE)</b> SrfI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Penghua Zhang  
Production Scientist  
18 Nov 2021



Michael Tonello  
Packaging Quality Control Inspector  
18 Nov 2021