

## New England Biolabs Certificate of Analysis

**Product Name:** PspOMI  
**Catalog Number:** R0653S  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10232263  
**Expiration Date:** 03/2026  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml rAlbumin (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0653S/L v2.0


PspOMI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0653SVIAL	PspOMI	10232252	Pass
B6004SVIAL	rCutSmart™ Buffer	10229454	Pass

Assay Name/Specification	Lot # 10232263
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 units of PspOMI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of PspOMI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of pXba DNA with PspOMI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PspOMI.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 200 units of PspOMI incubated for 16 hours at 37°C results in a DNA pattern free of	<b>Pass</b>

Assay Name/Specification	Lot # 10232263
detectable nuclease degradation as determined by agarose gel electrophoresis.	
<b>Protein Purity Assay (SDS-PAGE)</b> PspOMI is $\geq 95\%$ pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	<b>Pass</b>
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 20 units of PspOMI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is $\leq 1$ E. coli genome.	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Ana Egana  
Production Scientist  
21 Mar 2024



Michael Tonello  
Packaging Quality Control Inspector  
21 Mar 2024