New England BioLabs Inc.

be INSPIRED
drive DISCOVERY
stay GENUINE

240 County Road
Ipswich, MA 01938-2723

Tel 978-927-5054
Fax 978-921-1350

www.neb.com
info@neb.com

New England Biolabs
Certificate of Analysis

Product Name:	*BstAPI*
Catalog Number:	*R0654L*
Concentration:	*5,000 U/ml*
Unit Definition:	*One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 60°C in a total reaction volume of 50 µl.*
Packaging Lot Number:	*10060443*
Expiration Date:	*09/2021*
Storage Temperature:	*-20°C*
Storage Conditions:	*50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA*
Specification Version:	*PS-R0654S/L v1.0*

BstAPI Component List			
NEB Part Number	**Component Description**	**Lot Number**	**Individual QC Result**
R0654LVIAL	BstAPI	10055508	**Pass**
B7204SVIAL	CutSmart® Buffer	10053983	**Pass**

Assay Name/Specification	Lot # 10060443
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 25 units of BstAPI incubated for 4 hours at 60ºC releases <0.1% of the total radioactivity.	**Pass**
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BstAPI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16ºC. Of these ligated fragments, >95% can be recut with BstAPI.	**Pass**
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 5 Units of BstAPI incubated for 16 hours at 60ºC results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	**Pass**

This product has been tested and shown to be in compliance with all specifications.

_____

Jianying Luo
Production Scientist
20 Sep 2019

_____

Michael Tonello
Packaging Quality Control Inspector
25 Nov 2019

R0654L / Lot: 10060443