

## New England Biolabs Certificate of Analysis


**Product Name:** NheI-HF<sup>®</sup>  
**Catalog Number:** R3131M  
**Concentration:** 100,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot Number:** 10055443  
**Expiration Date:** 09/2021  
**Storage Temperature:** -20°C  
**Storage Conditions:** 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton X-100, 200 µg/ml BSA  
**Specification Version:** PS-R3131M v1.0

NheI-HF <sup>®</sup> Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3131M VIAL	NheI-HF <sup>®</sup>	10055444	Pass
B7204S VIAL	CutSmart <sup>®</sup> Buffer	10050275	Pass
B7024S VIAL	Gel Loading Dye, Purple (6X)	10043911	Pass

Assay Name/Specification	Lot # 10055443
<b>Blue-White Screening (Terminal Integrity)</b> A sample of LITMUS28i vector linearized with a 10-fold excess of NheI-HF <sup>™</sup> , religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 Units of NheI-HF <sup>™</sup> incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 300 units of NheI-HF <sup>™</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 100-fold over-digestion of Lambda HindIII DNA with NheI-HF <sup>™</sup> , >95% of the DNA	Pass

Assay Name/Specification	Lot # 10055443
<p>fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with NheI-HF™.</p> <p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda HindIII DNA and a minimum of 200 Units of NheI-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.



David Guo  
Production Scientist  
13 Aug 2019



Josh Hersey  
Packaging Quality Control Inspector  
19 Sep 2019