

## New England Biolabs Product Specification

<i>Product Name:</i>	<i>Standard Taq (Mg-free) Reaction Buffer Pack</i>
<i>Catalog #:</i>	<i>B9015S</i>
<i>Concentration:</i>	<i>10X Concentrate</i>
<i>Shelf Life:</i>	<i>60 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Composition (1X):</i>	<i>10 mM Tris-HCl, 50 mM KCl, (pH 8.3 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-B9015S v2.0</i>
<i>Effective Date:</i>	<i>12 Feb 2020</i>

### Assay Name/Specification (minimum release criteria)

**Endonuclease Activity (Nicking, Mg-Free Buffer)** - A 50 µl reaction in 2X Standard *Taq* (Mg-free) Reaction Buffer and 3 mM MgCl<sub>2</sub> containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Non-Specific DNase Activity (16 hour, Mg-Free Buffer)** - A 50 µl reaction in 2X Standard *Taq* (Mg-free) Reaction Buffer and 3 mM MgCl<sub>2</sub> containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**PCR Amplification (5 kb Lambda DNA, Mg-Free Buffer)** - A 50 µl reaction in Standard *Taq* (Mg-free) Reaction Buffer and 1.5 mM MgCl<sub>2</sub> in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 1.25 units of *Taq* DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product.

**pH (buffers/solutions)** - The pH of 10X Standard *Taq* (Mg-free) Reaction Buffer is between pH 8.2 and 8.4 at 25°C.

**Phosphatase Activity (pNPP, Buffer)** - A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl<sub>2</sub> containing 2.5 mM *p*-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl Standard *Taq* (Mg-free) Reaction Buffer incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

**qPCR DNA Contamination (*E. coli* Genomic, Buffer)** - A minimum of 1 µl of Standard *Taq* (Mg-free) Reaction Buffer is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.



---

## New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)
---

RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Standard <i>Taq</i> (Mg-free) Reaction Buffer is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.
--

One or more products referenced in this document may be covered by a 3rd-party trademark.  
Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Date 12 Feb 2020

---

Derek Robinson  
Director, Quality Control

