

## New England Biolabs Product Specification

<i>Product Name:</i>	<i>Q5<sup>®</sup> Hot Start High-Fidelity 2X Master Mix</i>
<i>Catalog #:</i>	<i>M0494S/L</i>
<i>Concentration:</i>	<i>2X</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Composition (1X):</i>	<i>Proprietary</i>
<i>Specification Version:</i>	<i>PS-M0494S/L v1.0</i>
<i>Effective Date:</i>	<i>06 Sep 2016</i>

### Assay Name/Specification (minimum release criteria)

**Endonuclease Activity (Nicking, Polymerase, dNTP)** - A 50 µl reaction in NEBuffer 2 in the presence of 400 µM dNTPs containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 units of Q5<sup>®</sup> High-Fidelity DNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Non-Specific DNase Activity (16 hour, Buffer)** - A 50 µl reaction in 1X Q5<sup>®</sup> Hot Start High-Fidelity Master Mix containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**PCR Amplification (20 kb Lambda DNA, Master Mix)** - A 50 µl reaction in 1X Q5<sup>®</sup> Hot Start High-Fidelity Master Mix and 1.0 µM primers containing 10 ng Lambda DNA for 22 cycles of PCR amplification results in the expected 20 kb product.

**PCR Amplification (7 kb Human Genomic DNA, Master Mix)** - A 50 µl reaction in 1X Q5<sup>®</sup> Hot Start High-Fidelity Master Mix and 0.5 µM primers containing 20 ng Human Genomic DNA for 30 cycles of PCR amplification results in the expected 7 kb product.

**PCR Amplification (Hot Start, Human Genomic DNA, Master Mix)** - A 25 µl reaction in 1X Q5<sup>®</sup> Hot Start High-Fidelity Master Mix and 0.5 µM primers containing 50 ng Human Genomic DNA for 25 cycles of PCR amplification results in the expected 665 bp product and a decrease in non-specific genomic bands after pre-incubation at room temperature for 1 hour, when compared to a non-hot start control reaction.

**Phosphatase Activity (pNPP)** - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl<sub>2</sub> containing 2.5 mM *p*-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Q5<sup>®</sup> High-Fidelity DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

**Protein Purity Assay (SDS-PAGE)** - Q5<sup>®</sup> High-Fidelity DNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.



---

## New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)
---

<p><b>qPCR DNA Contamination (<i>E. coli</i> Genomic)</b> - A minimum of 2 units of Q5<sup>®</sup> High-Fidelity DNA Polymerase is screened for the presence of <i>E. coli</i> genomic DNA using SYBR<sup>®</sup> Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is <math>\leq 1</math> <i>E. coli</i> genome.</p>
--

<p><b>RNase Activity (Extended Digestion)</b> - A 10 <math>\mu</math>l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 <math>\mu</math>l of Q5<sup>®</sup> Hot Start High-Fidelity 2X Master Mix is incubated at 37°C. After incubation for 4 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>
---



Date 06 Sep 2016

---

Derek Robinson  
Director of Quality Control

