

## New England Biolabs Product Specification

<b>Product Name:</b>	<i>PstI-HF</i> <sup>®</sup>
<b>Catalog #:</b>	R3140T/M
<b>Concentration:</b>	100,000 units/ml
<b>Unit Definition:</b>	One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.
<b>Shelf Life:</b>	24 months
<b>Storage Temp:</b>	-20°C
<b>Storage Conditions:</b>	250 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.15 % Triton X-100, 200 µg/ml rAlbumin, (pH 7.4 @ 25°C)
<b>Specification Version:</b>	PS-R3140T/M v3.0
<b>Effective Date:</b>	19 Feb 2024

### Assay Name/Specification (minimum release criteria)

**Exonuclease Activity (Radioactivity Release)** - A 50 µl reaction in rCutSmart<sup>™</sup> Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 200 units of PstI-HF<sup>®</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

**Ligation and Recutting (Terminal Integrity)** - After a 100-fold over-digestion of Lambda DNA with PstI-HF<sup>®</sup>, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PstI-HF<sup>®</sup>.

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in rCutSmart<sup>™</sup> Buffer containing 1 µg of Lambda DNA and a minimum of 200 units of PstI-HF<sup>®</sup> incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Protein Purity Assay (SDS-PAGE)** - PstI-HF<sup>®</sup> is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

**qPCR DNA Contamination (*E. coli* Genomic)** - A minimum of 20 units of PstI-HF<sup>®</sup> is screened for the presence of *E. coli* genomic DNA using SYBR<sup>®</sup> Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.





240 County Road  
Ipswich, MA 01938-2723

Tel 978-927-5054  
Fax 978-921-1350

www.neb.com  
info@neb.com

---

## New England Biolabs Product Specification

One or more products referenced in this document may be covered by a 3rd-party trademark.  
Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.

*Nancy Considine*

Date 19 Feb 2024

---

Nancy Considine  
Quality Approver

