

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	NEBuffer™ r2.1
Catalog Number:	B6002S
Concentration:	10 X Concentrate
Packaging Lot Number:	10243201
Expiration Date:	03/2027
Storage Temperature:	-20°C
Specification Version:	PS-B6002S v1.0
Composition (1X):	10 mM Tris-HCl, 50 mM NaCl, 10 mM MgCl2, 100 μg/ml rAlbumin, (pH 7.9 @ 25°C)

NEBuffer™ r2.1 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B6002SVIAL	NEBuffer™ r2.1	10231025	Pass	

Assay Name/Specification	Lot # 10243201
Conductivity (buffers/solutions) The conductivity of 10X NEBuffer™ r2.1 is between 55 and 62 mS/cm at 25°C.	Pass
Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 1X NEBuffer™ r2.1 containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X NEBuffer™ r2.1 containing 1 µg of Lambda DNA and 1 unit of Sphl incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X NEBuffer™ r2.1 containing 1 µg of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X NEBuffer™ r2.1 containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass





240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10243201
RNase Activity (Buffer) A 10 µl reaction in 1X NEBuffer™ r2.1 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
pH (buffers/solutions) The pH of 10X NEBuffer™ r2.1 is between pH 7.8 and 8.0 at 25°C.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

9 hore

Scott Schallehn Production Scientist 26 Mar 2024

Michae m. 11

Michael Tonello Packaging Quality Control Inspector 12 Jun 2024

