

New England Biolabs Certificate of Analysis

Product Name: Q5® Reaction Buffer Pack
Catalog Number: B9027S
Concentration: 5 X Concentrate
Packaging Lot Number: 10245730
Expiration Date: 12/2026
Storage Temperature: -20°C
Specification Version: PS-B9027S v2.0
Composition (1X): Proprietary

Q5® Reaction Buffer Pack Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B9028AVIAL	Q5® High GC Enhancer	10237049	Pass
B9027SVIAL	Q5® Reaction Buffer Pack	10239527	Pass

Assay Name/Specification	Lot # 10245730
Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 2X Q5® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 2X Q5® Reaction Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (20 kb Lambda DNA, Buffer) A 50 µl reaction in Q5® Reaction Buffer in the presence of 200 µM dNTPs and 1 µM primers containing 10 ng Lambda DNA with 1 unit of Q5® High-Fidelity DNA Polymerase for 22 cycles of PCR amplification results in the expected 20 kb product.	Pass
PCR Amplification (7 kb Human Genomic DNA, Buffer) A 50 µl reaction in Q5® Reaction Buffer in the presence of 200 µM dNTPs and 0.5 µM primers containing 20 ng Human Genomic DNA with 1 unit of Q5® High-Fidelity DNA Polymerase for 30 cycles of PCR amplification results in the expected 7 kb product.	Pass
Phosphatase Activity (pNPP, Buffer)	Pass

Assay Name/Specification	Lot # 10245730
<p>A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 80 µl Q5® Reaction Buffer incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	
<p>RNAse Activity Assay (4 Hour Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Q5® Reaction Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic, Buffer) A minimum of 1 µl of Q5® Reaction Buffer is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Trinh Nguyen
Production Scientist
15 Apr 2024



Michael Tonello
Packaging Quality Control Inspector
11 Jun 2024