

New England Biolabs Certificate of Analysis

Product Name: SplintR® Ligase

Catalog Number: M0375L
Concentration: 25,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme needed to ligate (to 50%

completion) 2 picomoles of a tripartite FAM-labeled DNA:RNA hybrid substrate in a 20 µl reaction at 25°C in 15 minutes in 1X SplintR®

Ligase Reaction Buffer.

Packaging Lot Number: 10241269
Expiration Date: 01/2026
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCI, 300 mM NaCI, 1 mM DTT, 0.1 mM EDTA, 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0375S/L v1.0

| SplintR® Ligase Component List | | | | |
|--------------------------------|-------------------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| M0375LVIAL | SplintR® Ligase | 10226193 | Pass | |
| B0375SVIAL | 10X SplintR® Ligase Reaction Buffer | 10226194 | Pass | |

| Assay Name/Specification | Lot # 10241269 |
|--|----------------|
| Endonuclease Activity (Nicking) | Pass |
| A 50 μl reaction in SplintR® Ligase Reaction Buffer containing 1 μg of supercoiled | |
| PhiX174 DNA and a minimum of 125 units of SplintR® Ligase incubated for 4 hours at | |
| 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | |
| Exonuclease Activity (Radioactivity Release) | Pass |
| A 50 μl reaction in SplintR® Ligase Reaction Buffer containing 1 μg of a mixture of | |
| single and double-stranded [³H] E. coli DNA and a minimum of 125 units of SplintR® | |
| Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | |
| Protein Purity Assay (SDS-PAGE) | Pass |
| SplintR® Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie | |
| Blue detection. | |
| RNase Activity (Extended Digestion) | Pass |
| 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA | |



M0375L / Lot: 10241269

Page 1 of 2



| Assay Name/Specification | Lot # 10241269 |
|---|----------------|
| and a minimum of 25 units of SplintR® Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel | |
| electrophoresis using fluorescent detection. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Production Scientist

06 Feb 2024

Josh Hersey

Packaging Quality Control Inspector

15 May 2024

