

New England Biolabs Certificate of Analysis

Product Name: EnGen® Spy Cas9 Nickase
Catalog Number: M0650T
Concentration: 20 µM
Unit Definition: N/A
Packaging Lot Number: 10243963
Expiration Date: 05/2026
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0650T v1.0

EnGen® Spy Cas9 Nickase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0650TVIAL	EnGen® Spy Cas9 Nickase	10241676	Pass
B6003SVIAL	NEBuffer™ r3.1	10227734	Pass

Assay Name/Specification	Lot # 10243963
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 1 pmol of EnGen® Spy Cas9 Nickase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 1 pmol of EnGen® Spy Cas9 Nickase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (EnGen® Spy Cas9 Nickase, Targeted Nicking) A 20 µl reaction in 1X NEBuffer 3.1 containing 20 nM 100 bp FAM and ROX labeled double stranded target DNA, 100 nM sgRNA and 100 nM EnGen® Spy Cas9 Nickase incubated for 1 hour at 37°C results in ≥90% nicking of the substrate DNA as determined by capillary electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1 pmol of EnGen® Spy Cas9 Nickase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel	Pass

Assay Name/Specification	Lot # 10243963
electrophoresis.	
Protein Purity Assay (SDS-PAGE) EnGen® Spy Cas9 Nickase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 pmol of EnGen® Spy Cas9 Nickase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Jessica Cane
Production Scientist
15 May 2024



Michael Tonello
Packaging Quality Control Inspector
03 Jun 2024