

New England Biolabs Certificate of Analysis

Product Name: EnGen[®] Spy dCas9 (SNAP-tag)
Catalog Number: M0652T
Concentration: 20 μ M
Unit Definition: N/A
Packaging Lot Number: 10153219
Expiration Date: 05/2024
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0652T v1.0

EnGen [®] Spy dCas9 (SNAP-tag) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0652TVIAL	EnGen [®] Spy dCas9 (SNAP-tag [®])	10153218	Pass
B6003SVIAL	NEBuffer [™] r3.1	10146824	Pass

Assay Name/Specification	Lot # 10153219
Protein Purity Assay (SDS-PAGE) EnGen [®] Spy dCas9 (SNAP-tag [®]) is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 pmol of EnGen [®] Spy dCas9 (SNAP-tag [®]) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Functional Testing (EnGen[®] Spy dCas9 (SNAP-tag[®]), Gel Shift Assay) A 20 μ l reaction in 1X NEBuffer 3.1 containing 20 nM 100 bp FAM labeled double stranded target DNA, 20 nM TAMRA-labeled off target DNA, 100 nM sgRNA and 100 nM EnGen [®] Spy dCas9 (SNAP-tag [®]) incubated for 15 minutes at 37°C results in \geq 90% binding of the substrate DNA as determined by electrophoretic mobility shift assay.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μ l reaction in NEBuffer 3.1 containing 1 μ g of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 1 pmol of EnGen [®] Spy dCas9 (SNAP-tag [®]) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

Assay Name/Specification	Lot # 10153219
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1 pmol of EnGen[®] Spy dCas9 (SNAP-tag[®]) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 1 pmol of EnGen[®] Spy dCas9 (SNAP-tag[®]) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Bhairavi Jani
Production Scientist
10 Jun 2022



Erin Varney
Packaging Quality Control Inspector
10 Jun 2022