

## New England Biolabs Certificate of Analysis

**Product Name:**  $\alpha$ 1-3,4,6 Galactosidase  
**Catalog Number:** P0747L  
**Concentration:** 8,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to cleave > 95% of the terminal,  $\alpha$ -D-galactose from 1 nmol Gal $\alpha$ 1-3Gal $\beta$ 1-4Gal-7-amino-4-methyl-coumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10  $\mu$ l.  
**Packaging Lot Number:** 10106406  
**Expiration Date:** 04/2022  
**Storage Temperature:** 4°C  
**Storage Conditions:** 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)  
**Specification Version:** PS-P0747S/L v1.0

<b><math>\alpha</math>1-3,4,6 Galactosidase Component List</b>			
<b>NEB Part Number</b>	<b>Component Description</b>	<b>Lot Number</b>	<b>Individual QC Result</b>
P0747LVIAL	$\alpha$ 1-3,4,6 Galactosidase	10106407	<b>Pass</b>
B9001SVIAL	Purified BSA	10098897	<b>Pass</b>
B1727SVIAL	10X GlycoBuffer 1	10092862	<b>Pass</b>

<b>Assay Name/Specification</b>	<b>Lot # 10106406</b>
<b>Glycosidase Activity (<math>\beta</math>-Mannosidase)</b> A 10 $\mu$ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled $\beta$ -Mannosidase substrate (Man $\beta$ 1-4Man $\beta$ 1-4Man-AMC) and 16 units of $\alpha$ 1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (<math>\beta</math>1-4 Galactosidase)</b> A 10 $\mu$ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled $\beta$ -Galactosidase substrate (Gal $\beta$ 1-4GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc -AMC) and 16 units of $\alpha$ 1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (<math>\beta</math>-N-Acetylglucosaminidase)</b> A 10 $\mu$ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled $\beta$ -N-Acetylglucosaminidase substrate (GlcNAc $\beta$ 1-4GlcNAc $\beta$ 1-4GlcNAc-AMC) and 16 units of $\alpha$ 1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>

Assay Name/Specification	Lot # 10106406
<p><b>Glycosidase Activity (<math>\beta</math>-N-Acetylgalactosaminidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\beta</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 16 units of <math>\alpha</math>1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-N-Acetylgalactosaminidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\alpha</math>1-3(Fuc<math>\alpha</math>1-2)Gal<math>\beta</math>1-4Glc-AMC) and 16 units of <math>\alpha</math>1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-6 Mannosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-6Man<math>\alpha</math>1-6(Man<math>\alpha</math>1-3)Man-AMC) and 16 units of <math>\alpha</math>1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-Glucosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Glucosidase substrate (Glc<math>\alpha</math>1-6Glc<math>\alpha</math>1-4Glc-AMC) and 16 units of <math>\alpha</math>1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\beta</math>1-3 Galactosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\beta</math>-Galactosidase substrate (Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 16 units of <math>\alpha</math>1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-Neuraminidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Neuraminidase substrate (Neu5Ac<math>\alpha</math>2-3Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 16 units of <math>\alpha</math>1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\beta</math>-Xylosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\beta</math>-Xylosidase substrate (Xyl<math>\beta</math>1-4Xyl<math>\beta</math>1-4Xyl<math>\beta</math>1-4Xyl-AMC) and 16 units of <math>\alpha</math>1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>

Assay Name/Specification	Lot # 10106406
<p><b>Glycosidase Activity (Endo F1, F2, H)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 16 units of α-1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (PNGase F)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 16 units of α-1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (Endo F2, F3)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 16 units of α-1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b> α-1-3, 4, 6 Galactosidase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (α1-3 Fucosidase)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 16 units of α-1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (α1-3 Mannosidase)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-3Manβ1-4GlcNAc-AMC) and 16 units of α-1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Protease Activity (SDS-PAGE)</b> A 20 µl reaction in 1X Glyco Buffer 1 containing 24 µg of a standard mixture of proteins and a minimum of 40 units of α-1-3, 4, 6 Galactosidase incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (α1-2 Fucosidase)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-2Galβ1-4Glc-AMC) and 16 units of α-1-3, 4, 6</p>	<b>Pass</b>

Assay Name/Specification	Lot # 10106406
Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Alicia Bielik  
Production Scientist  
15 Apr 2021



Michael Tonello  
Packaging Quality Control Inspector  
15 Apr 2021