

New England Biolabs Certificate of Analysis

Product Name: *Hpy188I*
Catalog Number: *R0617L*
Concentration: *10,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of pBR322 in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10243519*
Expiration Date: *04/2026*
Storage Temperature: *-20°C*
Storage Conditions: *10 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25C)*
Specification Version: *PS-R0617S/L v2.0*


Hpy188I Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0617LVIAL	Hpy188I	10235531	Pass
B6004SVIAL	rCutSmart™ Buffer	10238052	Pass

Assay Name/Specification	Lot # 10243519
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 30 units of Hpy188I incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of pBR322 DNA with Hpy188I, ~50% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with Hpy188I.	Pass
Non-Specific DNase Activity (16 hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of 10 units of Hpy188I incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.	Pass

Assay Name/Specification	Lot # 10243519
<p>Protein Purity Assay (SDS-PAGE) Hpy188I is $\geq 95\%$ pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of Hpy188I is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Ana Egana
Production Scientist
10 Jun 2024



Michael Tonello
Packaging Quality Control Inspector
10 Jun 2024