

## New England Biolabs Certificate of Analysis

**Product Name:** Agel-HF®  
**Catalog Number:** R3552L  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction of 50 µl.  
**Packaging Lot Number:** 10241912  
**Expiration Date:** 02/2026  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)  
**Specification Version:** PS-R3552S/L v2.0

### Agel-HF® Component List


NEB Part Number	Component Description	Lot Number	Individual QC Result
R3552LVIAL	Agel-HF®	10225347	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10236422	Pass
B6004SVIAL	rCutSmart™ Buffer	10238052	Pass

Assay Name/Specification	Lot # 10241912
<b>Blue-White Screening (Terminal Integrity)</b> A sample of LITMUS28i vector linearized with a 10-fold excess of Agel-HF®, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of Agel-HF® incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of Agel-HF® incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda DNA with Agel-HF®, >95% of the DNA	Pass

Assay Name/Specification	Lot # 10241912
<p>fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with Agel-HF®.</p>	
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of Agel-HF® incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b> Agel-HF® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 20 units of Agel-HF® is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Ana Egana  
Production Scientist  
04 Jun 2024



Michael Tonello  
Packaging Quality Control Inspector  
04 Jun 2024