

New England Biolabs Product Specification

Product Name: T4 Polynucleotide Kinase Reaction Buffer
Catalog #: B0201S
Concentration: 10X Concentrate
Shelf Life: 36 months
Storage Temp: -20°C
Composition (1X): 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT, (pH 7.6 @ 25°C)
Specification Version: PS-B0201S v2.0
Effective Date: 29 May 2024

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking, Buffer) - A 50 µl reaction in 1X T4 Polynucleotide Kinase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Functional Testing (Reaction Buffer) - Each lot of T4 Polynucleotide Kinase Reaction Buffer is functionally tested according to the unit definition for T4 Polynucleotide Kinase. Performance is directly compared to a control comprising a previously approved lot.

Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in 1X T4 Polynucleotide Kinase Reaction Buffer containing 1 µg of HaeIII digested PhiX174 RF I DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

RNase Activity (Buffer) - A 10 µl reaction in 1X T4 Polynucleotide Kinase Reaction Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by polyacrylamide gel electrophoresis.

One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.



Date 29 May 2024

Lauren Brown
Quality Approver

