240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350

www.neb.com info@neb.com

## **New England Biolabs Product Specification**

Product Name: Oligo  $d(T)_{25}$  Magnetic Beads

Catalog #: S1419S Concentration: 5 mg/ml Shelf Life: 36 months Storage Temp: 4°C

Storage Conditions: 0.05 % NaN3, 0.05 % Tween®20, 1 X PBS, (pH 7.4 @, 25°C)

Specification Version: PS-S1419S v3.0 Effective Date: 20 Dec 2023

## Assay Name/Specification (minimum release criteria)

Binding Capacity (Magnetic Beads) - Oligo  $d(T)_{25}$  Magnetic Beads ( 500  $\mu$ g ) were equilibrated and incubated with 100  $\mu$ l of 67 μg/ml rA30 for 15 minutes at 25°C, then washed and the rA30 eluted. Binding capacity was determined to be >5 μg of rA30 per mg of beads.

Functional Testing (mRNA Isolation) - Oligo d(T)<sub>25</sub> Magnetic Beads were equilibrated and incubated with freshly prepared eukaryotic cell lysate for direct mRNA isolation. The beads were washed and the mRNA eluted. The eluate was evaluated on an Agilent Bioanalyzer and the enriched poly(A)+ RNA contains  $\leq 15\%$  rRNA.

Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in Oligo d(T)<sub>25</sub> Magnetic Bead Storage Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

RNase Activity (Extended Digestion) - A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of Oligo d(T)<sub>25</sub> Magnetic Beads is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lauren Brown Quality Approver







wer Brown

Date 20 Dec 2023